



NEWS RELEASE

Jan7KEYTRUDA® (pembrolizumab) Receives New Draft Recommendation from National Institute for Health and Care Excellence (NICE) in U.K. for First-Line Treatment for Advanced Melanoma

10/9/2015

KEYTRUDA is Currently Approved in 39 Countries, including the United States and throughout the European Union

Merck (NYSE:MRK), known as MSD outside the United States and Canada, today announced the National Institute for Health and Care Excellence (NICE) of the United Kingdom (U.K.) has issued a draft recommendation, in the form of a Final Appraisal Determination (FAD), recommending KEYTRUDA® (pembrolizumab) as a first-line treatment option for adults with advanced melanoma.

“We are pleased that the U.K. government has recognized the value of KEYTRUDA, and thank the government for its efforts to ensure that patients in the U.K. who have advanced melanoma have access to KEYTRUDA as soon as possible,” said Deepak Khanna, senior vice president and regional president, Europe, MSD Oncology. “Merck has demonstrated our strong commitment to ensuring KEYTRUDA would be available as quickly as possible to patients in the U.K., and around the world. Since the first approval in the United States just more than a year ago, KEYTRUDA has been approved in 39 countries, including throughout the European Union.”

In addition to today’s NICE recommendation for KEYTRUDA as a first-line treatment option for adults with advanced melanoma, earlier this week NICE issued final guidance recommending KEYTRUDA for the treatment of advanced melanoma after disease progression with ipilimumab. KEYTRUDA was the first medicine available through the U.K.

Early Access to Medicines Scheme (EAMS), which was introduced in the U.K. in 2014 to help patients with life-threatening or seriously debilitating conditions benefit from promising, innovative treatments before a European license has been granted.

“The availability of KEYTRUDA for first-line use in patients will be welcomed by the melanoma community,” said Gillian Nuttall, Founder of Melanoma UK. “Advanced melanoma is a very difficult disease to treat effectively and this treatment will give hope to many. We are delighted that patients will be able to access this treatment on the National Health Services and congratulate NICE on their swift decision making.”

Securing approvals for KEYTRUDA globally is a key element of Merck’s efforts to ensure that the medicine is broadly available for eligible patients with advanced melanoma who are in need of new options around the world. To date, KEYTRUDA has been approved for the treatment of certain patients with advanced melanoma by regulatory authorities in the United States and the EU, as well as Australia, Canada, Israel, Macau, New Zealand, Peru, South Korea, Switzerland, and the United Arab Emirates (UAE). Additionally, the U.S. Food and Drug Administration (FDA) recently approved KEYTRUDA for certain patients with advanced non-small cell lung cancer ([link](#)).

About KEYTRUDA® (pembrolizumab) Injection 100 mg

KEYTRUDA is a humanized monoclonal antibody that works by increasing the ability of the body’s immune system to help detect and fight tumor cells. KEYTRUDA blocks the interaction between PD-1 and its ligands, PD-L1 and PD-L2, and may affect both tumor cells and healthy cells.

KEYTRUDA is indicated in the U.S. at a dose of 2 mg/kg administered as an intravenous infusion over 30 minutes every three weeks for the treatment of patients with unresectable or metastatic melanoma and disease progression following ipilimumab and, if BRAF V600 mutation positive, a BRAF inhibitor. KEYTRUDA is also indicated for the treatment of metastatic non-small cell lung cancer (NSCLC) whose tumors express PD-L1 as determined by an FDA-approved test with disease progression on or after platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving KEYTRUDA. In the U.S., these indications are approved under accelerated approval based on tumor response rate and durability of response; an improvement in survival or disease-related symptoms has not yet been established. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Selected Safety Information for KEYTRUDA in Advanced Melanoma Trial

Pneumonitis occurred in 12 (2.9%) of 411 patients, including Grade 2 or 3 cases in 8 (1.9%) and 1 (0.2%) patients, respectively, receiving KEYTRUDA. Monitor patients for signs and symptoms of pneumonitis. Evaluate suspected

pneumonitis with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 pneumonitis.

Colitis (including microscopic colitis) occurred in 4 (1%) of 411 patients, including Grade 2 or 3 cases in 1 (0.2%) and 2 (0.5%) patients, respectively, receiving KEYTRUDA. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold KEYTRUDA for Grade 2 or 3; permanently discontinue KEYTRUDA for Grade 4 colitis.

Hepatitis (including autoimmune hepatitis) occurred in 2 (0.5%) of 411 patients, including a Grade 4 case in 1 (0.2%) patient, receiving KEYTRUDA. Monitor patients for changes in liver function. Administer corticosteroids for Grade 2 or greater hepatitis and, based on severity of liver enzyme elevations, withhold or discontinue KEYTRUDA.

Hypophysitis occurred in 2 (0.5%) of 411 patients, including a Grade 2 case in 1 and a Grade 4 case in 1 (0.2% each) patient, receiving KEYTRUDA. Monitor patients for signs and symptoms of hypophysitis (including hypopituitarism and adrenal insufficiency). Administer corticosteroids for Grade 2 or greater hypophysitis. Withhold KEYTRUDA for Grade 2; withhold or discontinue for Grade 3; and permanently discontinue KEYTRUDA for Grade 4 hypophysitis.

Hyperthyroidism occurred in 5 (1.2%) of 411 patients, including Grade 2 or 3 cases in 2 (0.5%) and 1 (0.2%) patients, respectively, receiving KEYTRUDA. Hypothyroidism occurred in 34 (8.3%) of 411 patients, including a Grade 3 case in 1 (0.2%) patient, receiving KEYTRUDA. Thyroid disorders can occur at any time during treatment. Monitor patients for changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and for clinical signs and symptoms of thyroid disorders. Administer corticosteroids for Grade 3 or greater hyperthyroidism. Withhold KEYTRUDA for Grade 3; permanently discontinue KEYTRUDA for Grade 4 hyperthyroidism. Isolated hypothyroidism may be managed with replacement therapy without treatment interruption and without corticosteroids.

Type 1 diabetes mellitus, including diabetic ketoacidosis, has occurred in patients receiving KEYTRUDA. Monitor patients for hyperglycemia and other signs and symptoms of diabetes. Administer insulin for type 1 diabetes, and withhold KEYTRUDA in cases of severe hyperglycemia until metabolic control is achieved.

Nephritis occurred in 3 (0.7%) patients, consisting of one case of Grade 2 autoimmune nephritis (0.2%) and two cases of interstitial nephritis with renal failure (0.5%), one Grade 3 and one Grade 4. Monitor patients for changes in renal function. Administer corticosteroids for Grade 2 or greater nephritis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 nephritis.

Other clinically important immune-mediated adverse reactions can occur. The following clinically significant immune-mediated adverse reactions occurred in patients treated with KEYTRUDA: exfoliative dermatitis, uveitis,

arthritis, myositis, pancreatitis, hemolytic anemia, partial seizures arising in a patient with inflammatory foci in brain parenchyma, severe dermatitis including bullous pemphigoid, myasthenic syndrome, optic neuritis, and rhabdomyolysis.

For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm etiology or exclude other causes. Based on the severity of the adverse reaction, withhold KEYTRUDA and administer corticosteroids. Upon improvement of the adverse reaction to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Restart KEYTRUDA if the adverse reaction remains at Grade 1 or less. Permanently discontinue KEYTRUDA for any severe or Grade 3 immune-mediated adverse reaction that recurs and for any life-threatening immune-mediated adverse reaction.

Infusion-related reactions, including severe and life-threatening reactions, have occurred in patients receiving KEYTRUDA. Monitor patients for signs and symptoms of infusion-related reactions including rigors, chills, wheezing, pruritus, flushing, rash, hypotension, hypoxemia, and fever. For severe or life-threatening reactions, stop infusion and permanently discontinue KEYTRUDA.

Based on its mechanism of action, KEYTRUDA may cause fetal harm when administered to a pregnant woman. If used during pregnancy, or if the patient becomes pregnant during treatment, apprise the patient of the potential hazard to a fetus. Advise females of reproductive potential to use highly effective contraception during treatment and for 4 months after the last dose of KEYTRUDA.

KEYTRUDA was discontinued for adverse reactions in 9% of 411 patients. Adverse reactions, reported in at least two patients, that led to discontinuation of KEYTRUDA were: pneumonitis, renal failure, and pain. Serious adverse reactions occurred in 36% of patients. The most frequent serious adverse reactions, reported in 2% or more of patients, were renal failure, dyspnea, pneumonia, and cellulitis.

The most common adverse reactions (reported in at least 20% of patients) were fatigue (47%), cough (30%), nausea (30%), pruritus (30%), rash (29%), decreased appetite (26%), constipation (21%), arthralgia (20%), and diarrhea (20%).

The recommended dose of KEYTRUDA is 2 mg/kg administered as an intravenous infusion over 30 minutes every three weeks until disease progression or unacceptable toxicity. No formal pharmacokinetic drug interaction studies have been conducted with KEYTRUDA. It is not known whether KEYTRUDA is excreted in human milk. Because many drugs are excreted in human milk, instruct women to discontinue nursing during treatment with KEYTRUDA. Safety and effectiveness of KEYTRUDA have not been established in pediatric patients.

Selected Safety Information for KEYTRUDA in Metastatic NSCLC Trial

Pneumonitis occurred in 19 (3.5%) of 550 patients, including Grade 2 (1.1%), 3 (1.3%), 4 (0.4%), or 5 (0.2%) pneumonitis in patients receiving KEYTRUDA. Monitor patients for signs and symptoms of pneumonitis. Evaluate suspected pneumonitis with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 or recurrent Grade 2 pneumonitis.

Colitis occurred in 4 (0.7 %) of 550 patients, including Grade 2 (0.2%) or 3 (0.4%) colitis in patients receiving KEYTRUDA. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold KEYTRUDA for Grade 2 or 3; permanently discontinue KEYTRUDA for Grade 4 colitis.

Hepatitis occurred in patients receiving KEYTRUDA. Monitor patients for changes in liver function. Administer corticosteroids for Grade 2 or greater hepatitis and, based on severity of liver enzyme elevations, withhold or discontinue KEYTRUDA.

Hypophysitis occurred in 1 (0.2 %) of 550 patients, which was Grade 3 in severity. Monitor patients for signs and symptoms of hypophysitis (including hypopituitarism and adrenal insufficiency). Administer corticosteroids and hormone replacement as indicated. Withhold KEYTRUDA for Grade 2 and withhold or discontinue for Grade 3 or Grade 4 hypophysitis.

Hyperthyroidism occurred in 10 (1.8%) of 550 patients, including Grade 2 (0.7%) or 3 (0.3%). Hypothyroidism occurred in 38 (6.9%) of 550 patients, including Grade 2 (5.5%) or 3 (0.2%). Thyroid disorders can occur at any time during treatment. Monitor patients for changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and for clinical signs and symptoms of thyroid disorders. Administer replacement hormones for hypothyroidism and manage hyperthyroidism with thionamides and beta-blockers as appropriate. Withhold or discontinue KEYTRUDA for Grade 3 or Grade 4 hyperthyroidism.

Type 1 diabetes mellitus, including diabetic ketoacidosis, has occurred in patients receiving KEYTRUDA. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Administer insulin for type 1 diabetes, and withhold KEYTRUDA and administer anti-hyperglycemics in patients with severe hyperglycemia.

Nephritis occurred in patients receiving KEYTRUDA. Monitor patients for changes in renal function. Administer corticosteroids for Grade 2 or greater nephritis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 nephritis.

For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm etiology or exclude other causes. Based on the severity of the adverse reaction, withhold KEYTRUDA and administer corticosteroids. Upon improvement of the adverse reaction to Grade 1 or less, initiate corticosteroid taper and continue to taper

over at least 1 month. Resume KEYTRUDA when the adverse reaction remains at Grade 1 or less following steroid taper. Permanently discontinue KEYTRUDA for any severe or Grade 3 immune-mediated adverse reaction that recurs and for any life-threatening immune-mediated adverse reaction.

The following clinically significant, immune-mediated adverse reactions occurred in patients treated with KEYTRUDA: rash, vasculitis, hemolytic anemia, serum sickness, myasthenia gravis, bullous pemphigoid, and Guillain-Barré syndrome.

Infusion-related reactions, including severe and life-threatening reactions, have occurred in patients receiving KEYTRUDA. Monitor patients for signs and symptoms of infusion-related reactions including rigors, chills, wheezing, pruritus, flushing, rash, hypotension, hypoxemia, and fever. For severe or life-threatening reactions, stop infusion and permanently discontinue KEYTRUDA.

Based on its mechanism of action, KEYTRUDA can cause fetal harm when administered to a pregnant woman. If used during pregnancy, or if the patient becomes pregnant during treatment, apprise the patient of the potential hazard to a fetus. Advise females of reproductive potential to use highly effective contraception during treatment and for 4 months after the last dose of KEYTRUDA.

KEYTRUDA was discontinued due to adverse reactions in 14% of patients. Serious adverse reactions occurred in 38% of patients. The most frequent serious adverse reactions reported in 2% or more of patients were pleural effusion, pneumonia, dyspnea, pulmonary embolism, and pneumonitis.

The most common adverse reactions (reported in at least 20% of patients) were fatigue (44%), decreased appetite (25%), dyspnea (23%), and cough (29%).

Our Focus on Cancer

Our goal is to translate breakthrough science into innovative oncology medicines to help people with cancer worldwide. At Merck Oncology, helping people fight cancer is our passion and supporting accessibility to our cancer medicines is our commitment. Our focus is on pursuing research in immuno-oncology and we are accelerating every step in the journey – from lab to clinic – to potentially bring new hope to people with cancer. For more information about our oncology clinical trials, visit www.merck.com/clinicaltrials.

About Merck

Today's Merck is a global healthcare leader working to help the world be well. Merck is known as MSD outside the United States and Canada. Through our prescription medicines, vaccines, biologic therapies and animal health

products, we work with customers and operate in more than 140 countries to deliver innovative health solutions. We also demonstrate our commitment to increasing access to healthcare through far-reaching policies, programs and partnerships. For more information, visit www.merck.com and connect with us on **Twitter**, **Facebook** and **YouTube**.

Forward-Looking Statement of Merck & Co., Inc., Kenilworth, N.J., USA

This news release of Merck & Co., Inc., Kenilworth, NJ, USA (the “company”) includes “forward-looking statements” within the meaning of the safe harbor provisions of the United States Private Securities Litigation Reform Act of 1995. These statements are based upon the current beliefs and expectations of the company’s management and are subject to significant risks and uncertainties. There can be no guarantees with respect to pipeline products that the products will receive the necessary regulatory approvals or that they will prove to be commercially successful. If underlying assumptions prove inaccurate or risks or uncertainties materialize, actual results may differ materially from those set forth in the forward-looking statements.

Risks and uncertainties include, but are not limited to, general industry conditions and competition; general economic factors, including interest rate and currency exchange rate fluctuations; the impact of pharmaceutical industry regulation and healthcare legislation in the United States and internationally; global trends toward healthcare cost containment; technological advances, new products and patents attained by competitors; challenges inherent in new product development, including obtaining regulatory approval; the company’s ability to accurately predict future market conditions; manufacturing difficulties or delays; financial instability of international economies and sovereign risk; dependence on the effectiveness of the company’s patents and other protections for innovative products; and the exposure to litigation, including patent litigation, and/or regulatory actions.

The company undertakes no obligation to publicly update any forward-looking statement, whether as a result of new information, future events or otherwise. Additional factors that could cause results to differ materially from those described in the forward-looking statements can be found in the company’s 2014 Annual Report on Form 10-K and the company’s other filings with the Securities and Exchange Commission (SEC) available at the SEC’s Internet site (www.sec.gov).

###

Please see Prescribing Information for KEYTRUDA (pembrolizumab) at http://www.merck.com/product/usa/pi_circulars/k/keytruda/keytruda_pi.pdf and the Medication Guide for KEYTRUDA at http://www.merck.com/product/usa/pi_circulars/k/keytruda/keytruda_mg.pdf

MerckMedia:Pamela Eisele, 267-305-3558orAn Phan, 908-236-1130orInvestors:Teri Loxam, 908-740-1986orJustin Holko, 908-740-1879

text